# **Retraction Announcement**

This original article has been retracted on January 26th, 2023 on the request of the authors because of dispute regarding copyright of data use. - *Editor* 

*Retraction in:* Ma T, Zhang S, Huang D, Zhang G, Chen B, Zhang N. Endoscopic-assisted lateral neck dissection and open lateral neck dissection in the treatment of lateral neck lymph node metastasis in papillary thyroid carcinoma: A comparison of therapeutic effect. Pak J Med Sci. 2022;38(7):1905-1910. doi: https://doi.org/10.12669/pjms.38.7.5826

**Original Article** 

Open Access

Endoscopic-assisted lateral neck dissection and open lateral neck dissection in the treatment of lateral neck lymph node metastasis in papillary thyroid carcinoma: A comparison of therapeutic effect

Tao Ma<sup>1</sup>, Shuai Zhang<sup>2</sup>, Dongmei Huang<sup>3</sup>, Gang Zhang<sup>4</sup>, Boyi Chen<sup>5</sup>, Ning ang<sup>6</sup>

# ABSTRACT

**Objectives:** To compare the therapeutic effect of endoscopic-assisted lateral new dissection and open lateral neck dissection in the treatment of lateral neck lymph node metastasis of public with papillary thyroid carcinoma (PTC).

**Methods:** A retrospective analysis was carried out focusing on the general time t data of 86 patients with PTC treated Baoding No.1 Central Hospital from January 2020 to Septement 2021. According to different surgical methods, enrolled patients were divided into the endotropic surgement (n = 34) and the open surgery group (n = 52). Further comparison was performed on the operation indexes [operation time, postoperative length of stay in hospital, number of dissected lymp modes (central area, lateral cervical area), number of metastatic lymph nodes (central area, lateral cervical area), hospitalization cost], postoperative complications, postoperative neck pain, neuronal discomfort score, and satisfaction with postoperative cosmetic effect.

**Results:** The operation time and hospitalization cost of e en scopic surgery group were higher than those of the open surgery group, and the intraoperative bloc s was lower than that of the open surgery group, with statistically significant differences ( 5). There was no significant difference in the length of stay in the hospital, the number of dissected syn ades, the number of metastatic lymph nodes and the detection rate of lymph nodes in zone II between the two groups (p>0.05). Furthermore, a statistically significant difference was observed in the incide of postoperative complications between the two groups, which was lower in the endoce fic surgery group (29.4%) than that in the open surgery group (51.9%) (p<0.05). There was no significate difference in postoperative neck pain scores between the two groups (p>0.05). While the pottonerative why numbness discomfort score, and satisfaction score with postoperative cosmetic effect e endoscopic surgery group were better than those in the open surgery group, and the difference was stat sally significant (p<0.05).

**Conclusion:** Endoscopic-assisted latent neck dissection can reduce intraoperative blood loss and postoperative complication bidence in the treatment of lateral neck lymph node metastasis of PTC. However, it has the discover again of longer operation time and high hospitalization costs.

; Open lateral neck dissection; Papillary thyroid carcinoma; Lymph node

## doi: https://doi.org/10.12669/pjms.38.7.5826

## How to cite

metastasis

**KEYWORDS:** 

Ma T, Zhang S, New D, Zhang G, Chen B, Zhang N. Endoscopic-assisted lateral neck dissection and open lateral neck dissection in the treatment of the real neck lymph node metastasis in papillary thyroid carcinoma: A comparison of the rapeutic effect. Pak J Med Sci. 2022;38(7):1905-1910. doi: https://doi.org/10.12669/pjms.38.7.5826

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Correspondence: Shuai Zhang, Email: zs180701@163.com

*	Received for Publication:	December 5, 2021
*	Revision Received:	May 16, 2022
*	Corrected & Edited:	May 21, 2022
*	Revision Accepted: *	May 28, 2022

oscopic-

# INTRODUCTION

Thyroid cancer is the most common malignant tumor of the endocrine system, and its incidence in females is significantly higher than that in males.<sup>1</sup> Papillary thyroid carcinoma (PTC) comprises almost 80% of all thyroid cancers.<sup>2</sup> Traditional

open surgery is the gold standard for the treatment of PTC, and its 10-year survival rate reaches 90% after treatment.3 However, during lymph node dissection in the lateral neck in the process of traditional open surgery, it is necessary to make "L"-shaped, reverse "L"- or long "collar"-shaped incision in the patient's neck.<sup>4</sup> It may produce neck scars that cannot be covered, which will have an adverse physical and psychological impact on the patients. With the development of the concept of minimally invasive surgery, endoscopy in surgery is becoming more widely available in the clinical setting. Endoscopic thyroid surgery includes total endoscopic thyroid surgery and endoscopicassisted thyroid surgery. Among them, endoscopicassisted thyroid surgery can expose the anatomical structure of cervical nerves and blood vessels, and has the advantages of simple operation, beginnerfriendly, small separation wound, small incision and short learning curve, which has a wider application in clinical practice. The safety and cosmetic effect are the primary concerns of current clinical research on endoscopic and open surgeries in the treatment of PTC.<sup>5</sup> The present research was a retrospective study to analyze the surgical indexes postoperative complications, postoperative n pain, neck numbness discomfort score, and he satisfaction of postoperative cosmetic effect. It i expected to better understand the advantages and disadvantages of both surgeries and ovide a potential reference for clinical practice.

# METHODS

A retrospective analysis was carrie focusing on the general clinical of 86 patients with N 1 Central Hospital PTC treated in Baodi mb 2021. According from January 2020 to S ical me. enrolled patients to different were divi a o endoscopic surgery group curgery group (n = 52). In the (n = 34)d op with an average age of  $(31.4\pm7.4)$ endoscop and 31 fema years old (19 years); The type of operation was total thyroidectomy in 12 cases and unilateral lobectomy in 22 cases, including five cases on the left, 10 cases on the right and 19 cases on both sides. While in the open surgery group, there were 10 males and 42 females, with an average age of  $(33.9\pm7.3)$  years old  $(20\sim58 \text{ years})$ ; The operation type was total thyroidectomy in 17 cases and unilateral lobectomy in 35 cases, including 13 cases on the left, 23 cases on the right and 16 cases on both sides

*Ethical Approval:* This study was approved by the Institutional Ethics Committee of Baoding No.1 Central Hospital on September 20, 2022 (No.2022033), and written informed consent was obtained from all participants.

## Inclusion criteria:

- Patients who met the diagnostic criteria of PTC in the Guidelines of the American Thyroid Association (ATA)<sup>6</sup>;
- The newly treated patients to had no previous history of neck radiother
- Patients with tumor demeter them and who had received lateral neck maps ode dissection within the selective period of stay in the hospital<sup>7</sup>;
- Patients with the paximum diameter of metastatic lym, and ≤2cm.

# Exclusioneria

- Patient with cancer cells spreading distally;
- Patients
  Mere malignant tumors;
- Defients who were converted from endoscopic surgery to open surgery;

Patien with lymph node fusion and tumor n of trachea, esophagus, nerve, etc.<sup>8</sup>

There was no significant difference in general data (gender, age, type of operation, dissection e) between the two groups (p>0.05). This study was approved by the Medical Ethics Committee of our hospital, and all patients had provided informed consent.

Open lateral neck dissection was performed in the open surgery group: After routine anesthesia, an arc-shaped incision of about 12 ~ 15 cm in length was made along the dermatoglyph at 1.5  $\sim 2.0$  cm above the anterior cervical sternal notch. The location of the incision was close to the side of the dissection. The skin and platysma were cut layer by layer, which was up to the level of the digastric muscle, down to the clavicle, and outside to the posterior edge of the sternocleidomastoid muscle. Pay attention to avoid cervical veins and nerve plexus. Patients in this group were provided with total thyroidectomy + central lymph node dissection or hemithyroidectomy and isthmectomy + central lymph node dissection on the affected side. Intraoperatively, the accessory nerve was dissociated and protected, the sternocleidomastoid muscle was dissociated, associated with the dissociation of lymph nodes and soft tissues in the lateral area of the internal jugular vein; meanwhile, attention was paid to expose and protect the greater auricular nerve and the lesser occipital nerve, cut and ligate the transverse carotid artery and vein,

and sharply separate the internal jugular vein from the surface of the internal jugular vein; the enlarged lymph nodes and soft tissues in zones II, III and IV were completely removed along the surface of deep cervical fascia; then, the venous angle was sutured carefully, followed by the dissociation of the soft tissue inside the internal jugular vein, exposure of the carotid sheath, exposure and protection of the internal jugular vein and vagus nerve, and removal the lymph nodes and soft tissue inside the internal jugular vein. Postoperatively, additional attention was attached to the examination of bleeding points, lymphatic leakage, as well as the integrity of nerves and vessels.9 The incision was sutured after catheterization for drainage and hemostasis.

Endoscopic assisted lateral neck dissection was adopted in the endoscopic surgery group:<sup>10</sup> The incision was made at 1.0 ~ 1.5 cm above the anterior cervical sternal notch, with a length of about 6cm. The thyroid and central lymph nodes were treated like that in the open surgery group. Endoscopic instruments were placed into the anterior cervical incision. The layers of incision were the same as those in the open surgery group, followed by the construction of the operation space and determination of the range of operation. Intraoperatively, accessory nerve was dissociated and protected using an ultrasonic knife, the sternocleidomastoic muscle was dissociated, and the lymph 1 les and soft tissues were also dissociated in the ral area of the internal jugular vein; a further ration

was performed to expose and protect the greater auricular nerve and the lesser occipital nerve, cut and ligate the transverse carotid artery and vein, and separate the internal jugular vein; the enlarged lymph nodes and soft tissues in zones II, III and IV were completely removed along the surface of deep cervical fascia; and the venous angle was sutured carefully, followed by the dissociation of the soft tissue inside the internal juchar vein, exposure of the carotid sheath, exposure protection of the internal jugular vein and vag e, and removal the lymph nodes and soft t ue in. the internal with the bleeding jugular vein. Postoperative point, lymphatic leakage, the grity of nerves the incision was sutured and vessels confir after catheterization for arginage and hemostasis. era n time, postoperative Surgical indexes: length on the pital, number of dissected lymph n (central area, lateral cervical area), number of hereitic lymph nodes (central area, lateral cervical a...a), and hospitalization cost. *coperative complications:*<sup>11</sup> Hoarseness, hy-P

arath pidism, hypocalcemia and postoperaom ti oma.

Postoperative neck pain, neck numbness discomfort score, and satisfaction with a postoperacosmetic effect: All scores were evaluated by ubjective evaluation method. Three days after the operation, according to the presence of pain, the scores of pain were: one point, no pain; two points, mild pain; three points, moderate pain;

	Endoscopic surgery group (n=34)	Open surgery group (n=52)	t/Z	р
Operation time (min)	172.6±44.0	137.2±31.9	4.329	0.001
Blood loss (	47.7±10.1	60.5±13.4	4.775	0.001
Postoper ve ler h of stay in hospital (d)	6.0(4.0, 7.0)	6.0(5.0, 6.8)	0.673	0.501
Number Nis n nodes (n)				
Central area	10.5(8.0, 15.0)	9.0(5.0, 15.0)	1.762	0.078
Lateral cervical a c	27.0(22.0, 36.5)	27.5(18.3, 37.5)	0.588	0.557
Number of metastatic lymph nodes (n)				
Central area	4.0(1.8, 7.3)	3.5(2.0, 6.0)	0.382	0.703
Lateral cervical area	4.0(1.0, 6.0)	3.0(1.0, 4.0)	1.033	0.302
Number of lymph nodes in zone II (n)	10.0(5.8, 15.5)	7.0(4.0, 11.8)	1.841	0.066
Number of metastatic lymph nodes in zone II (n)	0(0, 1.0)	0(0, 1.0)	0.363	0.717
Detection rate of lymph nodes in zone II (%)	0(0, 0.1)	0(0, 0.2)	0.769	0.442
Hospitalization cost (million yuan)	4.3±0.5	3.7±0.4	6.157	0.001

Table-I: parison of surgical indexes between the two groups.

p

#### Tao Ma et al.

	Endoscopic surgery group (n=34)	Open surgery group (n=52)	$\chi^2$	p	
 Hoarseness (n%)	5(14.7)	8(15.4)	0.007	0.932	
Hypoparathyroidism (n%)	6(17.6)	11(21.2)	0.159	0.690	
Hypocalcemia (n%)	5(14.7)	9(17.3)	0.102	0.749	
Postoperative hematoma (n%)	0	6(11.5)	4.2	0.040	
 Total (n%)	10(29.4)	27(51.9)	4.2	039	

Table-II: Comparison of postoperative complications between the two groups.

four points, severe pain; and five points, extremely severe pain.<sup>12</sup> Neck numbness discomfort was evaluated one month after operation (one point: no discomfort, two points: mild discomfort, three points: moderate discomfort, four points: severe discomfort and five points: extremely severe discomfort).<sup>13</sup> In addition, at three months after the operation, the cosmetic effect satisfaction score was given, including one point of very satisfied, two points of satisfied, three points of generally satisfied, 4 points of dissatisfied and five points of very dissatisfied.<sup>14</sup>

Statistical Analysis: SPSS 26.0 statistical software was used for statistical analysis. The measurem data conforming to the normal distribution v expressed by mean±standard deviation (Mean±SD) and the t-test was used for the comparison between groups. While the measurement date hat did not conform to the normal distribution were expressed by median and inter le spacin, and comparison between groups wa lized by using the Mann-Whitney U test. The count ata were expressed by rate and corresponded between groups using the  $\chi^2$  test. P<0.07 mean that the difference was statistically signific

# RESUL

As shown in while L, the operation time and hospitalize the ost of L, endoscopic surgery group were higher on those of the open surgery group, and the intraoperative blood loss was lower than that of the open surgery group, with statistically significant differences (p<0.05). Lole-I. There was no significant difference in the length of stay in the hospital, the number of dissected lymph nodes, the number of monstatic lymph nodes and the detection of lym<sub>1</sub> – nodes in zone II between the two groups (p>0.05).

A statistical prioriticant difference was observed in the incidence of postoperative complications be seen the two groups, which was lower in the enoscope surgery group than that in the open surgery open (p<0.05).Table-II. Besides, hoarseness and hypoparathyroidism occurred temporarily after the operation.

There was no significant difference in postoperative neck pain scores between the two groups (p>0.05). While the postoperative neck numbness discomfort score, and satisfaction score with postoperative cosmetic effect in the endoscopic surgery group were better than those in the open surgery group, and the difference was statistically significant (p<0.05). Corresponding results are shown in Table-III.

## DISCUSSION

PTC has a high risk of lymph node metastasis, with an incidence of about 30% ~ 80%.<sup>15</sup> Traditional open surgery is a safe and effective choice for the treatment of lateral neck lymph node metastasis, while the formed scars after operation may disturb the normal daily life of patients.<sup>16</sup> With the progress

Table-III: Comparison of scores between the two groups.

	Endoscopic surgery group (n=34)	Open surgery group (n=52)	Ζ	р
Neck pain score (point)	2(1, 2)	2(1, 2)	0.472	0.637
Neck numbness discomfort score (point)	1(1, 2)	2(1, 2)	1.987	0.047
Satisfaction score with postoperative cosmetic effect (point)	2(1, 2)	3(2, 3)	5.036	0.001

of the concept of biology-medicine-society, there is an extensive application of minimally invasive endoscopic surgery in clinic. Minimally invasive endoscopic cleaning of PTC refers to the process of establishing operation space through hook suspension technology via minimal neck incision, and completing the operation with the use of a 5 mm endoscope and ultrasonic knife.<sup>17</sup> It has no visual dead angle during operation in the neck, and has significant advantages in the surgery of thyroid cancer. According to prior research,<sup>18</sup> despite no significant difference between endoscopic surgery and traditional open surgery in efficacy and safety, the former surgery can significantly reduce the length of neck incision and has a cosmetic effect to some extent; and its cosmetic effect is superior to the latter one considering a lower position of the incision that is easy to cover. However, endoscopic surgery is a novel surgical approach that has not been fully popularized; and there are few physicians at home and abroad who acquire proficiency in applying this technique for the treatment of thyroid cancer, resulting in controversy in its safety for treating tumors.19

As discovered in our study, there was significant difference in the number of disse lymph nodes, the number of metastatic lymph nodes and the detection rate of lymph noder in zone II between the two groups (p>0.05). It ggested that endoscopic dissection has a similar the eutic , both of Aich effect with traditional open su can play a satisfied role in lympk. dissection. In the aspect of safety, although there obvious difference in postoperative sk pain score between ide e of postoperative groups (p>0.05), the i r pic urgery group was complications in the end lower than the op ery group (29.4% vs. 51.9%) panied by better postoperative 200 neck nu ness scomfort score and satisfaction score with cosmetic effect than those up (p<0.05). These results support in the open that endoscopic regery is superior to open surgery in safety and can reduce the discomfort caused by surgery. In terms of potential causes, the use of endoscope intraoperatively can broaden the operation field of vision to accelerate the discovery of micro-nerves and blood vessels, resulting in the reduced risk of nerve and blood vessel injury during the operation;<sup>20</sup> secondly, the application of endoscope offers flexible lighting that can change in intensity and lens distance, which may contribute to observing the tissue condition;<sup>21</sup> and thirdly,

endoscopic surgery has a small incision, which can meet the patients' requirements for cosmetic effect; simultaneously, rather than traditional ligation, electrocoagulation is used to stop bleeding, which can reduce intraoperative blood loss and postoperative hematoma infection risk.

However and importantly, endoscopic surgery still has some defects, such as longer operation time and high hospitalization cost As for the prolonged operative duration, the major of physicians are unfamiliar with relevant g ns considering a delayed popularization and lication of a. I this regard, endoscopic surgery in C the operation time of this supry is relatively traditional open surgery prolonged compar which has been vid y and skillfully operated by doctors. While has been reported that the theoretic eration of endoscopic surgery can be shorte after the mastering of this technique<sup>22</sup> since endose is surgery indeed has significant advantages in ... operation processes with great culty and high precision requirements. di

**The limit ons of the study** It has the disadvantages of here operation time and high hospitalization costs. Findings in our study require further erification based on larger sample size and longer ration of study owing to the smaller sample size and shorter follow-up period in the present study.

## CONCLUSIONS

Endoscopic-assisted lateral neck dissection can reduce intraoperative blood loss and postoperative complication incidence in the treatment of lateral neck lymph node metastasis of PTC.

### Conflic of interest: None

*Funding:* This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

#### REFERENCES

- Nan BY, Xiong GF, Zhao ZR, Gu X, Huang XS. Comprehensive Identification of Potential Crucial Genes and miRNA-mRNA Regulatory Networks in Papillary Thyroid Cancer. Biomed Res Int. 2021;2021:6752141. doi: 10.1155/2021/6752141
- McDow AD, Saucke MC, Marka NA, Long KL, Pitt SC. Thyroid Lobectomy for Low-Risk Papillary Thyroid Cancer: A National Survey of Low- and High-Volume Surgeons. Ann Surg Oncol. 2021;28(7):3568-3575. doi: 10.1245/s10434-021-09898-9
- McDow AD, Pitt SC. Extent of Surgery for Low-Risk Differentiated Thyroid Cancer. Surg Clin North Am. 2019;99(4):599-610. doi: 10.1016/j.suc.2019.04.003

- Park VY, Han K, Kim HJ, Lee E, Youk JH, Kim EK, et al. Radiomics signature for prediction of lateral lymph node metastasis in conventional papillary thyroid carcinoma. PLoS One. 2020;15(1):e0227315. doi: 10.1371/journal. pone.0227315
- Wang TS, Sosa JA. Thyroid surgery for differentiated thyroid cancer - recent advances and future directions. Nat Rev Endocrinol. 2018;14(11):670-683. doi: 10.1038/s41574-018-0080-7
- Koc AM, Adıbelli ZH, Erkul Z, Sahin Y, Dilek I. Comparison of diagnostic accuracy of ACR-TIRADS, American Thyroid Association (ATA), and EU-TIRADS guidelines in detecting thyroid malignancy. Eur J Radiol. 2020;133:109390. doi: 10.1016/j.ejrad.2020.109390
- Bae SY, Jung SP, Choe JH, Kim JS, Kim JH. Prediction of lateral neck lymph node metastasis according to preoperative calcitonin level and tumor size for medullary thyroid carcinoma. Kaohsiung J Med Sci. 2019;35(12):772-777. doi: 10.1002/kjm2.12122
- Abraham E, Roshan D, Tran B, Wykes J, Campbell P, Ebrahimi A. The extent of extrathyroidal extension is a key determinant of prognosis in T4a papillary thyroid cancer. J Surg Oncol. 2019;120(6):1016-1022. doi:10.1002/jso.25683
- Yazici D, Colakoglu B, Saglam B, Sezer H, Kapran Y, Aydin O, et al. Effect of prophylactic central neck dissection on the surgical outcomes in papillary thyroid cancer: experience in a single center. Eur Arch Otorhinolaryngol. 2020;277(5):1491-1497. doi: 10.1007/s00405-020-05830-1
- Van Le Q, Ngo DQ, Le DT, Ngo QX. ASO Author Reflections: A New Procedure for Modified Radical Neck Dissection via the Transoral Endoscopic Approach. Ann Surg Oncol. 2021;28(5):2767. doi: 10.1245/s10434-020-09515-1
- Burger F, Fritsch H, Zwierzina M, Prommegger R, Konschreide M. Postoperative Hypoparathyroidism in Thyroid Sure ry Anatomic-Surgical Mapping of the Parathyroids an Implications for Thyroid Surgery. Sci Rep. 2010 (1):15700 doi: 10.1038/s41598-019-52189-3
- Saxena S, Gonsette K, Terram W, Huybrecht, and Jahrwold DA, Cappello M, et al. Gradual withdrawal of reconstantial delays initial post-operative analgement and after a proid surgery; double-blinded, randomized patrolled trial. BMC Anesthesiol. 2019;19(1):60. doi: 10.1186/j.ue?71/919-0731-9
- Kim KL, Park GY, Kwon DR, Kwon DY, Ludak SG, Cho HK. Airway invasion in non-templogically ill patients with dysphagia: Contributing actors and associated problems during swallowing processive discretional study. Medicine (Balting 4, 2 0;99(45):e22977. doi: 10.1097/MP.000000022.
- 14. Lee DW ang Jeong JH, Kwak SG, Choi YY, Tae K. Cosme outcor a after transoral robotic thyroidectomy: Compound axillary, postauricular, and convents approaches. Oral Oncol. 2021;114:105139. doi: 10.1016/j.ok.applogy.2020.105139
- Feng JW, Qin Ye J, Pan H, Jiang Y, Qu Z. Predictive Factors for Lateral Lymph Node Metastasis and Skip Metastasis in Papillary Thyroid Carcinoma. Endocr Pathol. 2020;31(1):67-76. doi: 10.1007/s12022-019-09599-w
- Cihan A, Cihan E. Interrelation Between Appearance Anxiety and Sexual Functions in Women: The Role of Surgical Scars, Morphologic Features, and Accompanying Depression. J Sex Med. 2019;16(11):1769-1778. doi: 10.1016/j. jsxm.2019.08.004
- Zhang D, Park D, Sun H, Anuwong A, Tufano R, Kim HY, et al. Indications, benefits and risks of transoral thyroidectomy. Best Pract Res Clin Endocrinol Metab. 2019;33(4):101280. doi: 10.1016/j.beem.2019.05.004

- 18. Wongwattana P, Laoveerakul P, Santeerapharp A. A comparison of efficacy and quality of life between transoral endoscopic thyroidectomy vestibular approach (TOETVA) and endoscopic thyroidectomy axillo-breast approach (ETABA) in thyroid surgery: non-randomized clinical trial. Eur Arch Otorhinolaryngol. 2021;278(10):4043-4049. doi: 10.1007/s00405-021-06639-2
- Ma JJ, Zhang DB, Zhang WF, Wang X. Application of Nanocarbon in Breast Approach Endoscopic Thyroidectomy Thyroid Cancer Surgery. J Laparoendosc Adv Surg Tech A. 2020;30(5):547-552. doi: 10.1089/hap.2019.0794
- Ji YB, Ko SH, Song CM, Subara S, Lee BJ, Wu CW, et al. Feasibility and efficacy of traoperative neural monitoring in remote accession and endoscopic thyroidectomy. Oral Oncol. 2 2:103:10 doi: 10.1016/j. oraloncology.2020.104617
- Zhang D, Gao L, Xie L, He G, Chengen C, et al. Comparison Between Video-Assist L and Open Lateral Neck Dissection for Papillary Thyracce. Jorna with Lateral Neck Lymph Node Metastasis A Josephtive Randomized Study. J Laparoendosc Advance Ten A. 2017;27(11):1151-1157. doi: 10.108 (Jap.2016.06)
- 22. McHeurer R. Is Prophylactic Central Compartment Neck Dissection plicated for Clinically Node-Negative Papillary Thyroid Care of the Answer is Dependent on How the Data are Interplated and the Weight Given to the Risks Benefits. Ann Surg Oncol. 2018;25(11):3123-3124. doi: 10.1245-110434-018-6649-5

### Au contributions:

**TM & SZ:** Designed this study, prepared this enuscript, are responsible and accountable for he accuracy and integrity of the work.

**DH & GZ:** Collected and analyzed clinical data.

**BC & NZ:** Data analysis, Significantly revised this manuscript.

#### Authors:

- 1. Tao Ma,
- Shuai Zhang,
  Dongmei Hua
  - Dongmei Huang, Department of Thyroid and Neck Tumor, Tianjin Medical University Cancer Institute and Hospital, National Clinical Research Center for Cancer, Key Laboratory of Cancer Prevention and Therapy, Tianjin's Clinical Research Center for Cancer, Tianjin 300060, China.
- 4. Gang Zhang,
- 5. Boyi Chen,
- 6. Ning Zhang,
- 1,2,4-6: Department of General Surgery, Baoding No.1 Central Hospital, Baoding, 071000, Hebei, China.

*Note:* Tao Ma and Dongmei Huang are both considered as co-first authors

Correspondence:

Shuai Zhang, Department of General Surgery, Baoding No.1 Central Hospital, Baoding, 071000, Hebei, China. Email: zs180701@163.com